





Analysing intermolecular interactions 101 - Full Interaction Maps Try



CCDC
advancing structural science

Table of Contents

Try – it's your turn!.....	2
Objectives	2
Example 1. Using Full Interaction Maps to assess polymorph stability	3
Conclusions for Example 1	5
Example 2. Investigating the potential for co-crystallisation.....	6
Conclusions for Example 2	7
Example 3. Investigating a series of halogenated molecules.....	8
Conclusions for Example 3	10
Bonus Exercises (optional)	10
Conclusions.....	11
Next steps.....	11
Glossary	12



Try – it's your turn!

In the second part of the *Analysing intermolecular interactions 101 – Full Interaction Maps* module, “U Try”, you will review the concepts and features presented in the first part, “U Watch”, and try them out yourself. If at any time you want to rewatch the videos, you can find them from the course webpage (<https://www.ccdc.cam.ac.uk/Community/educationalresources/CSDU/Analysing-intermolecular-interactions-FIMs/>).

This tutorial will introduce you to the Full Interaction Maps feature included with Mercury under the CSD-Materials and CSD-Discovery toolsets.

Before beginning, ensure that you have Mercury installed on your computer and an active CSD license. For further information, please contact your site administrator or the CCDC ([Contact Us](https://www.ccdc.cam.ac.uk/theccdcprofile/contactus/) - <https://www.ccdc.cam.ac.uk/theccdcprofile/contactus/>).


Objectives

In this workshop you will:

- Learn how to produce Full Interaction Maps (FIMs) for a given molecule and how to interpret the results.
- See how the FIMs analysis can be used for co-crystal design.
- See how FIMs can be used to investigate the differences in the interaction preferences of a molecule when substitutions are made across a series.

This workshop will take approximately **40 minutes** to be completed.

Note: The words in *Blue Italic* in the text are reported in the [Glossary](#) at the end of this handout.




Example of Full Interaction Maps.


Example 1. Using Full Interaction Maps to assess polymorph stability

The stability of a given crystal structure is a balance between the intramolecular conformation and the intermolecular packing of the molecules in the crystalline state. One method for understanding the relative stability of crystal structures is to compare the observed intermolecular interactions with preferred geometries for that type of interaction.

This example will look at two known polymorphs of sulfathiazole to answer the question: **How do the interactions in each polymorph compare with what is expected?** You will learn how to produce a Full Interaction Map for a given structure and how to interpret this map.



Sulfathiazole, 2D representation and CSD entry SUTHAZ19

1. Launch Mercury by clicking its icon . In the Structure Navigator toolbar, type SUTHAZ19 to bring up the structure for the Form V polymorph of sulfathiazole.
2. Click on the *CSD-Materials* menu or *CSD-Discovery* menu and select *Full Interaction Maps...* Note: If the CSD-Materials menu bar is inactive, click on "Help" and "Activate CSD-Materials" before proceeding further.
3. In the *Full Interaction Maps* dialogue box, you will see several options. On the left you will find options to change the display contour levels. On the right, you will see a list of functional groups to be used as probes. For the purposes of this tutorial, we will keep the default options. These typically work well for most situations, but if you know you are looking for a specific functional group, or if you want to change the look of the map, you will want to change these settings.
4. Click the **Calculate Maps** button to start.

The screenshot illustrates the steps to generate a Full Interaction Map in Mercury.

1. The **Structure Navigator** window shows 'SUTHAZ19' selected in the 'Crystal Structures' list.


2. The **CSD-Materials** menu is open, and 'Full Interaction Maps...' is selected.

3. The **Full Interaction Maps** dialog box is shown. The 'Map Contour Levels' section has three checked options: 'Display first contour with initial level of 2.0', 'Display second contour with initial level of 4.0', and 'Display third contour with initial level of 6.0'. The 'Hotspots' section has 'Generate hotspots in the map' unchecked. The 'Probe' list on the right includes 'Uncharged NH Nitrogen' (checked), 'Charged NH Nitrogen', 'RNH3 Nitrogen', 'Alcohol Oxygen', 'Carbonyl Oxygen', 'Water Oxygen', 'Oxygen Atom', 'Methyl Carbon', 'Aromatic CH Carbon' (checked), 'C-F Fluorine', 'C-Cl Chlorine', 'C-Br Bromine', and 'C-I Iodine'. A color scale bar is visible on the right.


4. The 'Calculate Maps' button at the bottom of the dialog is highlighted.

5. The generated map will now be displayed in the main Mercury window. Notice the three different colours in the map. **Red** regions of the map denote areas in which there is a high probability of locating a hydrogen bond **acceptor**. **Blue** regions denote hydrogen bond **donors**, and **brown** regions indicate **hydrophobic** preferences. It can be easier to view FIMS with a white background. To change the background colour, right click in an empty space on the visualiser, select "Display Options" and set the Background to a single colour of white.
6. Now we want to see how the overall packing of this polymorph fits with the Full Interaction Map we have generated. Tick the box for *H-Bond* in the Display Options toolbar.
7. Double click the *H-Bond* line to launch the *Define H-bonds* dialogue. In this dialogue, tick the box for "Require hydrogen atom to be present". Click **OK** to apply the change.
8. Now you will see dashed **red** lines in the Mercury window that indicate where hydrogen bonding interactions/contacts are present.
9. Click on these contacts to generate nearby molecules. You will see that in each case, the interaction falls within the contour range for the expected type. This indicates that the packing of Form V satisfies the expected interaction landscape of this conformation of sulfathiazole.
10. Now let's look at the Form I polymorph. In the Structure Navigator toolbar, type SUTHAZ16 to load this structure.
11. Click the **Reset** button in the Display Options toolbar to remove all the hydrogen bonding interactions.
12. Note that $Z' = 2$ for Form I; meaning there are two unique molecules in the asymmetric unit. We will focus on one unique molecule in the Form I structure.


5




7




6




8




9




10



11



13




13. Tick the box to Label atoms in the Display Options toolbar. Locate which molecule contains N1. This is the molecule we will focus on. Once you have located it, remember which one it is, but untick the Label atoms box to keep the display clean.
14. Right-click on the molecule that contains N1 then go to "Selection > Select Molecule". This will select the entire molecule. Alternatively, hold the Shift key and click the molecule that contains N1.
15. If the *Full Interaction Maps* dialogue is still open, simply click **Calculate Maps** again. Otherwise, follow steps 2-4 to re-open the dialogue.
16. You should now have a Full Interaction Map surrounding molecule 1.
17. Following **Steps 6-9** above, turn on the hydrogen bonding interactions and click to expand the interaction around the N1 atom.
18. Notice that one of the three interactions (N1-H2...O2) falls well outside the predicted region for a hydrogen bond acceptor. This suggests that this interaction has a non-ideal geometry and is likely to be significantly less stabilising than the interactions in Form V.

Conclusions for Example 1

The observed polymorphs of sulfathiazole exhibit different interactions, but also noticeably different geometries for those interactions. We can use knowledge-based approaches to compare the observed intermolecular interactions in two polymorphs with the preferred geometries for these interaction types. Full Interaction Maps indicate that Form V (known to be the most stable of the sulfathiazole polymorphs) has interactions which are near to ideal, whereas Form I (known to be metastable) has non-ideal interactions.


You should now know how to generate a Full Interaction Map of a molecule through Mercury and how to explore hydrogen bonding interactions in a crystal structure in relation to the predicted interaction landscape.



Example 2. Investigating the potential for co-crystallisation

For this tutorial, we will investigate the molecule anastrozole. This compound is a non-steroidal aromatase inhibitor used to treat breast cancer. The only published crystal structure is of the pure form. The molecule also contains no conventional hydrogen bond donors.

In this example, you will generate a Full Interaction Map for this molecule and investigate the interaction landscape. You will see how this information can be used to design a co-crystal.



1. In the Mercury window, type SATHOL in the Structure Navigator toolbar to load the structure of anastrozole.
2. Click the *CSD-Materials* menu or *CSD-Discovery* menu and select *Full Interaction Maps...* from the dropdown menu.
3. Keep the default settings.
4. Click **Calculate Maps** to generate the Full Interaction Map for SATHOL.
5. The resulting map shows several areas for hydrogen bond donors (blue regions); two high-probability areas near the nitrogen atoms of the triazole ring, and two slightly lower probability regions near the cyano groups.
6. While there are no true hydrogen bond donors in this structure, the C-H groups in the triazole ring could interact with possible hydrogen bond acceptors (red regions).
7. There is also a small brown region indicating the possibility for a hydrophobic or π - π interaction.

1

Structure Navigator

SATHOL Find

Crystal Structures	Space
SATHOL	C2/c
SATHOM	C2/c
SATHON	P41
SATHUR	P-1
SATHUS	C2/c
SATHUT	P43

2

CSD-Materials CSD-Discovery CSD-Materials CSD-Discovery CSD Python API He

Search

Calculations

Polymorph Assessment

Co-Crystal Design

Full Interaction Maps...

Hydrate Analyser...

Solvate Analyser...

Aromatics Analyser...

Conformer Generation...

Launch DASH

3

Full Interaction Maps

Options Maps Hotspots Log Files

Map Contour Levels

☒ Display first contour with initial level of 2.0

☒ Display second contour with initial level of 4.0

☒ Display third contour with initial level of 6.0

Hotspots

☐ Generate hotspots in the map

Probe

☒ Uncharged NH Nitrogen

☐ Charged NH Nitrogen

☐ RNH3 Nitrogen

☐ Alcohol Oxygen

☐ Carbonyl Oxygen

☐ Water Oxygen

☐ Oxygen Atom

☐ Methyl Carbon

☒ Aromatic CH Carbon

☐ C-F Fluorine

☐ C-Cl Chlorine

☐ C-Br Bromine

☐ C-I Iodine

Colour

4

Calculate Maps Clear Maps & Hotspots Load Maps... Save Maps... Close

5

6







7

8. As there are no conventional hydrogen-bond donors in this structure, there will be nothing to display by ticking the H-Bond box in the Display Options toolbar. Instead, tick the Short Contact box.
9. Double click this bar to open the *Define Short Contacts* dialogue box. Change this so that we find contacts shorter than the sum of the vdW radii **plus** 0.15Å. Click **OK** to save your changes.
10. Let's first look at the region around the triazole ring. We can see that there are two C-H donors that will satisfy the interaction preferences of the two N acceptors in the triazole ring. Click on the two contacts (dashed red lines) to generate the neighbouring molecules.
11. Looking at the region around the cyano groups, there are some short contacts available for interactions, but these are not directed toward the high-probability areas.
12. From the acceptor probe maps (red contours), we can see that the main region of acceptor preference is only weakly satisfied by the cyano group from one molecule and the triazole ring from another. Click the contacts to expand nearby molecules. The other acceptor regions are not satisfied at all.
13. Finally, there is a weak π — π interaction between the two phenyl rings that matches with one of the hydrophobic regions.

Conclusions for Example 2

The Full Interaction Maps for SATHOL show that the highest probability regions for interactions, the donor regions for the triazole N atoms, are quite nicely satisfied. However, this comes at the cost of satisfying the interactions that would fulfill the acceptor sites. Since anastrozole has no classical H-bond donors of its own to fulfill those interactions, a molecule with one or more H-bond donors would be a good choice for a potential co-crystallisation agent.


There are no co-crystals or solvates of anastrozole yet found in the CSD.



Example 3. Investigating a series of halogenated molecules

When designing molecules to fit a specific purpose, chemists will often make subtle changes to a single functional group; for example, replacing fluorine with chlorine, bromine, or iodine. We can use Full Interaction Maps to investigate the differences in the interaction preferences of a molecule when these substitutions are made across a series.

This example will look at the series of halogenated benzoic acids to compare what differences there are across the series.







A series of halogenated benzoic acids

1. Launch Mercury and type PFBZAD15 in the Structure Navigator tool bar to bring up one of the parafluorobenzoic acid entries.
2. Click the *CSD-Materials* menu or *CSD-Discovery* menu and choose *Full Interaction Maps...* from the drop-down menu.
3. We will again use the default settings.
4. Click **Calculate Maps** to generate the Full Interaction Map for parafluorobenzoic acid.
5. Repeat **Steps 1-4** for the other molecules in this series. Open a new instance of Mercury each time so you can easily compare the results. Use refcodes CLBZAP10 for the Cl analogue, BRBZAP01 for the Br analogue, and BENMOW07 for the I analogue. These particular entries were chosen because they are relatively recent crystal structure determinations at low temperatures. To open multiple Mercury windows on a **Mac**, open a terminal window and run this command:

```
open -n -a "/Applications/CCDC/CSD_2020/mercury.app"
```

 This command tells the system to open a new instance (the "-n" flag) of an application (the "-a" flag) that is the given name in quotes.
6. Now let's compare the maps for each molecule in the series.

7. Look at the maps generated across the series. You should notice that as the size of the halogen increases across the series, there is an increased likelihood of finding an acceptor interacting with the halogen. This interaction is non-existent for the fluoro species, but well-defined for the bromo and iodo analogues.
8. Now we will investigate how this affects the packing in each crystal structure.
9. Start with the iodo analogue (BENMOW07). In the Display Options make sure the Short Contact settings are for "sum of vdW radii **plus** 0.15Å." If not, double click that line to open the Define Short Contacts dialog and change the settings. Click **OK** to accept these changes. Then tick the box to display Short Contacts.
10. With the short contacts displayed, you should see that most of the interactions are satisfied. The two high-probability donor and acceptor sites near the carboxylic acid group are satisfied, as are the weaker acceptor sites near the aromatic hydrogen atoms on the phenyl ring. The iodine from a neighbouring molecule serves as the acceptor for a halogen-halogen interaction.
11. Click the interactions highlighted in Figure 10 to expand these neighbouring molecules. You will notice that while the acceptor (red) contours on either side of the phenyl ring are roughly equivalent, only one region is satisfied well by an interaction. Iodine atoms from neighboring molecules fill this region on the opposite side.







12. Repeat **Steps 9-11** for the Br (BRBZAP01) and Cl (CLBZAP10) analogues. What similarities and differences do you see across the series so far? What do you expect to see for the fluoro analogue?
13. For the fluoro analogue (PFBZAD15), follow **Step 9** to generate the short contacts.
14. You will see that the interactions for the carboxylic acid group and for the two weak C-H donors on the phenyl ring are satisfied. However, there is no probability of an acceptor for the fluorine, and consequently there is only C and H in this region.
15. Click these contacts to expand the neighbouring molecules. Now you can see that there is no halogen-halogen interaction involving the fluorine.

Conclusions for Example 3

It is well understood that iodine forms halogen-halogen interactions more so than the other halogen atoms. Using Full Interaction Maps, we can clearly see how the probability of these interactions affects the overall packing in a crystal structure.

Bonus Exercises (optional)

- Try calculating Full Interaction Maps for the molecules in Example 3 using only the C-F and C-Cl probes. Do the results match your expectations?
- Use Full interaction maps for a crystal that exhibits jumping properties, L-pyroglutamic acid, before (LPYGLU07) and after a phase transition (LPYGLU08). What do you observe for the phase transition structure?
- Run a Full Interaction Maps calculation on your own molecules.




Conclusions

After this workshop you will:


- Know how to produce Full Interaction Maps (FIMs) for a given molecule and how to interpret the results.
- Have experience in exploring and assessing the hydrogen bonding interactions in a crystal structure in relation to the interaction landscape predicted by FIMs.
- Be familiar with how the FIMs analysis can be used for co-crystal design.
- Be familiar with how FIMs can be used to investigate the differences in the interaction preferences of a molecule when substitutions are made across a series.

Next steps


Now that you have completed the Try part of this CSDU module, you can head to the final part: the quiz! Go back to the module webpage (<https://www.ccdc.cam.ac.uk/Community/educationalresources/CSDU/Analysing-intermolecular-interactions-FIMs/>) and follow the instructions to complete the final test.



LPYGLU07



LPYGLU08



Glossary

Contour level

The number of times more than random an interaction is likely to occur in a specific region of space.

Hotspots


Hotspots represent the positions of highest local density for each contour Surface.

Van der Waals, Aromatic and Hydrogen Bond Interactions

Van der Waals forces are formed between atoms or molecules that are in each other's close proximity and are driven by induced electrical interaction. They are the weakest of all type of intermolecular attractions between molecules. However, with a lot of Van der Waals forces interacting between two molecules, the interaction can be very strong.

Aromatic Interactions are noncovalent interactions formed between aromatic rings. These interactions are important in material science since they will contribute to the overall crystal structure stability. The orientation of the aromatic ring can vary from parallel to T-shape, and we found during our DFT calculations that the T-shape interactions are very close in strength to the parallel displaced ones. Their strength is found between 0 and 16 kJ/mol based on DFT calculations.

Hydrogen Bonding occurs between donor-acceptor interactions precisely involving hydrogen atoms. The H-bonds interactions are classified as: strong (mostly covalent), moderate (mostly electrostatic) and weak (electrostatic). Their strength is observed to be between 12 and 30 kJ/mol.



<i>Interaction type</i>	<i>Strength (kJ/mol)</i>
<i>Van der Waals</i>	0.4-4.0
<i>Aromatics</i>	0-16
<i>Hydrogen Bonds</i>	12-30